



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

3. If the above method does not succeed, allow the small pieces of fresh tissue to remain exposed to the action of the air for ten to twenty minutes before placing in the Flemming solution. This should be at a temperature of 4° C. Time as in 2.
4. Wash in water about 24 hours.
5. Dehydrate by very gradual steps.
6. Clear from 95 per cent alcohol in cedar oil followed by xylol.
7. Imbed in paraffin.

QUINCE JELLY AS A CULTURE MEDIUM FOR EUGLENA

Turner (Anat. Rec., 12:3, April, 1917) develops a mode of culturing *Euglena* and other green protozoa in quince jelly, which has practical value for the general laboratory because of the ease of making and preserving the cultures. Cultures have been kept for more than a year, and may be successfully transplanted.

The methods and results may be summarized as follows:

1. Twenty grams of dry quince seed boiled for half an hour in 1.5 liters of distilled water produces enough of the exudate, after the seeds are strained out, to make a total culture mass of 2½ liters, by the addition of distilled water.
2. A still more fruitful medium was made by grinding the seeds to a fine powder or meal and leaving it in the jelly. Water to 2½ or 3 liters may be used with this amount of meal. This medium has the disadvantage of being opaque.
3. Cultures develop more rapidly in a slightly thinner medium, but more lasting cultures call for a thicker medium.
4. The medium should be made slightly alkaline, and the cultures kept at room temperature, in moderate light.
5. Test tubes, vials, and similar vessels furnishing large vertical range and little exposed surface are best.
6. The reactions of *Euglena* are modified in interesting ways, as compared with their usual behavior in water.

"VITAL DYES" IN TELEOSTS

Wislocki (Anat. Record. XII, 4, May 1917) reports on experiments with benzidine dyes injected into the peritoneal cavity of